

# Comunicaciones Breves

## SMALL MAMMALS IN PELLETS OF BIRDS OF PREY FROM A DESERT ECOTONE IN NORTHWESTERN PATAGONIA, ARGENTINA

PEQUEÑOS MAMÍFEROS EN EGAGRÓPILAS DE AVES RAPACES EN UN ECOTONO DESÉRTICO DEL NOROESTE DE PATAGONIA, ARGENTINA

Andrea del Pilar Tarquino-Carbonell<sup>1\*</sup>, Paola Turienzo<sup>2</sup> & José Manuel López<sup>1</sup>

<sup>1</sup>Instituto Argentino de Investigaciones de las Zonas Áridas (IADIZA), Centro Científico y Tecnológico, Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET). Avenida Ruiz Leal s/n Parque General San Martín, Mendoza, Argentina

<sup>2</sup>Universidad Nacional de Río Negro, Sede Andina, Bariloche, Argentina

\*atarquino@mendoza-conicet.gob.ar

**ABSTRACT.**- One of the distinctive features of the South American drylands is the presence of small mammals. This work presents the results of taphonomic and taxonomic analyses of micromammal ( $\leq 1$  kg) bones recovered from an isolated deposit in pellets of birds of prey found in a cave located in the Neuquén province (Argentina). Taphonomic analyses revealed a low proportion of elements modified by digestion and of broken elements, as well as a relatively high abundance of post-cranial skeletal elements, a pattern suggesting that pellets were ejected by barn owls (*Tyto furcata*). Remains from members of five rodent genera (*Calomys*, *Eligmodontia*, *Phyllotis*, *Oligmodontia*, *Ctenomys*) were identified. The results of these analyses should contribute to ongoing ecological studies, especially on birds of prey, of Northwestern Patagonia.

**KEYWORDS.**- *drylands, rodentia, skeletal remains, Strigiformes, taphonomy*

**RESUMEN.**- Uno de los rasgos distintivos de las tierras áridas sudamericanas es la composición de pequeños mamíferos. En este trabajo se presentan resultados tafonómicos y taxonómicos de restos óseos de micromamíferos ( $\leq 1$  kg) recuperados de egagrópilas de aves rapaces encontradas en una caverna localizada en la provincia de Neuquén (Argentina). El análisis tafonómico reveló bajas proporciones de elementos modificados por la digestión y de elementos rotos como también abundancias relativamente altas de elementos esqueléticos, un patrón que sugiere que las egagrópilas fueron acumuladas por lechuzas de campanario (*Tyto furcata*). Se identificaron restos de miembros de cinco géneros de roedores (*Calomys*, *Eligmodontia*, *Phyllotis*, *Oligmodontia*, *Ctenomys*). Los resultados de estos análisis deberían contribuir a los estudios ecológicos en curso, especialmente en aves de presa del noroeste de la Patagonia.

**PALABRAS CLAVE.**- *restos óseos, rodentia, Strigiformes, tafonomía, zonas áridas*

South America encompasses a diversity of arid habitats (i.e., drylands) occurring in tropical, highland, coastal, and continental biomes (Morello 1958). The small mammal fauna associated with these arid habitats varies widely with regard to taxonomic composition and species richness (Ojeda et al. 2000). One such biome, the Monte Desert, extends for over 2000 km in western Argentina, ranging from 24 to 4 degrees

(Brown and Pacheco 2006, Arana et al. 2021); this habitat, which covers an area of approximately 460 km<sup>2</sup>, resembles parts of the North American Sonoran and Chihuahuan deserts. Daily temperature ranges from 27 to 7°C (Cabrera 1971), with annual precipitation typically being < 100-450 mm (Ojeda and Tabeni 2009). A second arid biome, the Patagonian Steppe is the 8th largest desert in the world, with a surface area

of ca. 560 000 km<sup>2</sup> (Brown and Pacheco 2006, Murrie and Murrie 2010). Most of this cold desert, located in Argentina, receives less than 200 mm of precipitation per year (Paruelo et al. 1998), with daily temperatures ranging from 2 to 6°C in winter and 10 to 30°C in summer (Ojeda et al. 2000).

The small mammal communities in these arid habitats can differ markedly and these diverse landscapes are thought to be important factors in shaping the evolution of these communities (Ojeda et al. 2000). For example, although herbivores are the primary trophic level in both Monte Desert and Patagonian Steppe habitats, these two regions display marked differences in the associated assemblages of small mammals (Mares and Ojeda 1982). Importantly, arid ecosystems biomes have been characterized as having high levels of endemism of mammalian species (Mares 1992), suggesting that historical records of the small mammal communities in these habitats can be used as proxies to reconstruct paleoenvironmental conditions. Bird pellets are often a valuable source of modern and historical small mammal remains and thus provide an effective tool for inferring environmental conditions over multiple time scales (Pardiñas et al. 2012, Fernández et al. 2016a, Fernández et al. 2016b, Fernández and Pardiñas 2018, López et al. 2021b).

As part of efforts to understand the vertebrates communities in arid South American habitats, we analyzed bones and teeth from small mammals ( $\leq 1$  kg) recovered from modern bird pellets collected from Caverna del León, Neuquén Province, Argentina. The goals of these analyses were to identify the likely source (predator species depositing pellets) and to characterize the taxonomic composition and species richness of the small mammals present in this deposit. These data, in turn, are used to draw inferences

regarding environmental conditions in the vicinity of Caverna del León.

## METHODS


### Study area

Caverna del León (38°37'S, 70°13'W, 1310 m.a.s.l.) is located at the interface between Monte Desert and Patagonian Steppe habitats in southwestern Argentina (Figure 1). The cave is located 20 km from the town of Las Lajas, in the center of Neuquén Province. The cave and surrounding area are used for multiple recreational activities, including visits by various speleological groups (Martinez 2018).


The approximate length of the cave is 852 m, with a maximum height of 24 m. The cave is composed of two large rooms, known as the “big room” and the “lake room”. The entrance to the cave follows a 40° slope (i.e., a dejection cone) that opens into the big room; a sub horizontal conduit exits the far end of this space, providing access to the lake room (Figure 1b), which contains a body of water approximately 50 m in diameter and 40 m deep. Several smaller galleries lead from the lake room to other portions of the cave complex (Tambor or Las Calizas room); additional details of the cave structure are provided in Barredo et al. (2012).

### Data collection and analysis

Six bird pellets were recovered from the lake room of the cave complex during a speleological expedition conducted in December 2017. The predator depositing these pellets was not observed directly. The appearance of all pellets was consistent with those



**Figure 1.** Study area and Caverna del León map shows a) location of study site in southern south America bordering Monte Desert and Patagonian Steppe ecoregions (Brown and Pacheco 2006), and b) Cavern structure (Tarquino-Carbonell and Turienzo 2019).



**Figure 2.** Cranial and tooth elements of the species found in Caverna del León, focused on molar series showing a) right upper molars of *Phyllotis pehuenche*, b) right lower molars of *P. pehuenche*, c) left upper molars of *Eligmodontia* sp., d) right lower molars of *Eligmodontia* sp., e) left upper molars of *Calomys musculus*, f) left lower molars of *C. musculus*, g) right upper molars of *Oligoryzomys longicaudatus*, h) right lower molars of *Ctenomys* sp. Occlusal view. Scale: 1 mm.

of recently deposited samples. In the laboratory, the pellets were disaggregated with surgical instruments and all small mammal remains (teeth, bones) separated from other material contained in the pellets.

Taxonomic identifications were conducted by comparing small mammal remains with published descriptions and specimens housed in regional museum collections (CMI, Colección Mastozoológica IA-DIZA). Most such comparisons were based on cranial and dental material (Fernández et al. 2011, Patton et al. 2015), with materials isolated from pellets examined under a stereomicroscope (LEICA S6D; 6.3-40x) to facilitate identification of distinguishing characteristics. From these data, the number of identified specimens (NISP) and minimum number of individuals (MNI) were calculated (Lyman 2008). For taphonomic analyses, the relative abundance of skeletal (versus cranial) elements was examined, as were bone breakage patterns and evidence of digestive corrosion (Andrews 1990, Fernández et al. 2017). For each specimen, patterns of digestive corrosion on teeth (molars, incisors) and bones (femora, humeri) were classified as being light, moderate, heavy, or extreme based on the extent and intensity of the corrosion detected (Fernández et al. 2017, Montalvo et al. 2016).


To evaluate the relationship between the abundance of different skeletal elements in living mammals and their representation in our sample, two taphonomic indices were employed. The first index calculated the relationship between cranial and postcranial elements (postcranial/ cranial, femora + humeri / mandible + maxillae); the second index measured the relationship between distal and proximal parts of the skeleton (tibiae+ radii/ femora + humeri). For both indices, the completeness of individual bones was evaluated separately for cranial and postcranial remains (Andrews 1990), after which the percentage of complete bones was calculated for both cranial elements and post-cranial long bones (i.e., femora). A principal components analysis was then used to

determine which skeletal remains tended to cluster the relative abundances in our sample. Finally, tendencies of relative abundances were estimated. Both PCA and relative abundance analyses were conducted using the libraries stats and ggplot2 package in R (R Development Core Team 2020).


## RESULTS

Based on bone and dental remains, total NISP for our sample was 320, with a MNI of 8. Five species of rodents (Figure 2) were identified: *Calomys musculus* (NISP = 2), *Eligmodontia* sp. (NISP = 5), *Oligoryzomys longicaudatus* (NISP = 3), *Phyllotis pehuenche* (NISP = 6) and *Ctenomys* sp. (NISP = 2). Three of these taxa (*O. longicaudatus*, *Eligmodontia* sp., *P. pehuenche*) were each represented by a MNI= 2 (see Discussion). The remaining two taxa (*C. musculus*, *Ctenomys* sp.) were each represented by MNI = 1. Taphonomic analyses revealed that almost no diagnostic elements exhibited digestive corrosion (Table 1). The detected corrosion was typically characterized as light; only one element displayed moderate corrosion, with no examples of heavy or extreme digestion detected (Figure 3).

Principal component analysis included nocturnal and diurnal avian predators. Component 1 (54.33%) and 2 (22.53%) together accounted for 76.86% of the inertia (Figure 4). The hemimandibular/skull completeness and postcranial completeness were represented by percentages of 61.53% and 100%, respectively. The highest values of relative abundances of skeletal elements were represented by pelvis, tibiae and femora (Table 2). The pc/c index resulted considerable higher than 1 (3.31), indicating better representation of postcranial elements. The f+h/md+mx index (0.85) suggested a slightly better representation of cranial elements in comparison with postcranial ones. The value obtained for t+r/f+h index (1.18) showed a slightly better representation of distal parts in comparison to proximal elements of the skeleton.



**Figure 3.** Digestion process, light digestion on Cricetidae incisor. Scale: 1 mm.



**Figure 4.** Principal component analysis including values reported in other birds of prey according to Montalvo and Fernández (2019) compared with this study.


## DISCUSSION

This study reports on the taxonomic composition and species richness of a small mammal community at the interface between Monte Desert and Patagonian Steppe habitats, as revealed by analyses of recently deposited bird pellets. No taxa distinctive to Monte Desert habitats were detected. In contrast, two of the species identified (*O. longicaudatus*, *P. pehuenche*) are considered typical of Patagonian Steppe habitats. The presence of *P. pehuenche* in our sample is particularly intriguing; this recently described species (Jayat et al. 2021) is thought to be endemic to northwestern portions of the Patagonian Steppe biome. Identification of *P. pehuenche* in our sample was based on mor-

phological characters as well as geographic location. Morphologically, this species is distinguished by a skull that is not robust compared to other members of the genus, zygomatic arches that are comparatively not very expanded, and zygomatic notches that are narrower and less excavated than other species. The molar tooth row is longer but thinner than that in the *P. xanthopygus* complex. Additionally, the occlusal surface of the molars in *P. pehuenche* is more simplified, without the deep paraflexus on M2 or the mesoloph complex on M1 and/or M2 observed in *P. vaccarum* (Jayat et al. 2022). Geographically, *P. pehuenche* is the only member of this thought to occur in Neuquen Province. However, a recent taxonomic review revealed sympatry between this species and the *P. xanthopygus* complex in southern Mendoza, indicating that additional sampling in this region is needed to delimit more precisely the distributions of these species (Jayat et al. 2021, 2022).

## Taphonomic implications

Similar findings regarding the relative abundance of post-cranial skeletal elements and the frequency of digestive corrosion have been reported for analyses of small mammal remains obtained from barn owl pellets recovered from Monte Desert habitat (Figure 5). Owls generally swallow their prey whole, resulting in a high relative abundance of post-cranial skeletal elements and low frequencies of bone and tooth breakage (Andrews 1990). From a taphonomic perspective, predation by owls typically results in limited evidence of digestion of diagnostic elements (Andrews 1990). Barn owls have been implicated as the source of pellets containing small mammal remains in multiple studies from southern South America (Saavedra and Simonetti 1998, Pardiñas 1999, López and Chiavazza 2019, López 2020, Mignino et al. 2021), suggesting that barn owls are likely to be the primary source of pellets at sites from central western Argentina, including Caverna del León. Although other species of owls may also produce pellets containing small mammal remains, these species (i.e., *Athene cucularia*, *Bubo magellanicus*) tend to produce remains that display greater evidence of damage and digestion than was detected during this study (Montalvo et al. 2016, Montalvo and Fernández 2019, López et al. 2021b). In contrast, the high relative abundance of post-cranial skeletal elements in our sample resembles reports for other analyses of pellets produced by barn owls (Andrews 1990). Thus, the results of our taphonomic analyses suggest pellets at Caverna del León were deposited by members of the Strigiformes, with barn



**Figure 5.** Relative abundances of elements according to Andrews (1990), Fernández et al. (2016b) and López (2020) compared with this study. Scap: scapulae. Fem: femurs. Hum: humeri. Mand: Mandibles. Max: Maxillae. Pel: Pelvis. Rad: Radii. Tib: Tibiae. Uln: Ulnae.

**Table 1.** Number and percentages of diagnostic elements modified by digestive corrosion indicated as absent (A), light (L), moderated (M), heavy (H) and extreme (E).

	A	%	L	%	M	%	H	%	E	%
Incisors <i>in situ</i>	20	76.92	5	19.23	1	3.85	0	0	0	0
Isolated incisors	2	100	0	0	0	0	0	0	0	0
Molars <i>in situ</i>	67	93.05	5	6.94	0	0	0	0	0	0
Isolated molars	2	100	0	0	0	0	0	0	0	0
Femora	11	84.61	2	15.38	0	0	0	0	0	0
Humeri	8	88.89	1	11.11	0	0	0	0	0	0

**Table 2.** Minimal number of elements (MNE), relative abundances (RA), number of identified specimens (NISP), broken elements and breakage percentages.

Element	MNE	RA (%)	NISP	n broken	Breakage (%)
Maxillae	15	93.75	15	9	60
Mandible	11	68.75	11	1	9
Scapulae	10	62.5	10	7	70
Humerii	9	56.25	9	0	0
Ulnae	10	62.5	10	0	0
Radii	12	75	12	0	0
Pelvis	14	87.5	14	0	0
Femurs	13	81.25	13	0	0
Tibiae	14	87.5	14	0	0

owls (*Tyto furcata*) being a particularly likely contributor to our sample (López 2020).

Our conclusion that the pellets obtained from Caverna del León were deposited by owls is similar to findings from other sites located in the Monte Desert-Patagonian Steppe ecotone (Fernández et al. 2016b, López 2020). Accordingly, our findings add an important comparative data point to the growing number of studies that use bird pellets to characterize the small mammal communities at arid sites in South America. Although numerous studies have examined taphonomic patterns generated by different predators, differences have also been observed in the taphonomic trends generated by the same type of predator in different environmental regions, often associated with the size and availability of prey. Our findings have implications to strengthen environmental reconstructions in different taxonomic and taphonomic discussions.

#### ACKNOWLEDGEMENTS

To the members of the speleological group GEMA for collecting the material. To the members of the Entomology Laboratory of IADIZA, José Ramírez, for their assistance with our photographic stuff. Pablo Teta from “Bernardino Rivadavia” Natural Sciences Argentine Museum, for his collaboration in the taxonomic identifications. A. Zarco, J. Mignino and E. A. Lacey for their valuable comments in the manuscript. PICT 2019-01515 (Agencia Nacional de Promoción de la Investigación, el Desarrollo Tecnológico y la Innovación, Argentina) and SIIP 06/G805 (Universidad Nacional de Cuyo). We thank three anonymous reviewers for comments that improved our manuscript.

#### LITERATURE CITED

- ANDREWS P (1990) Small Mammal Taphonomy. Pp. 487-494 in: Lindsay EH, Fahlbusch V y Mein P, (eds) European Neogene Mammal Chronology. NATO Science Series A: (NSSA, volume 180), Springer, New York
- ARANA MD, NATALE E, FERRETTI N, ROMANO G, OGGERO A, MARTÍNEZ G, POSADAS P AND MORRONE JJ (2021) Esquema biogeográfico de la República Argentina. *Opera lilloana* 56, Fundación Miguel Lillo, Tucumán
- BARREDO S, GABRIELE N, GARRIDO A AND REDONTE G (2012) Los principales sistemas de cavernas cársticas de la provincia del Neuquén. *Revista de la Asociación Geológica Argentina* 69:556-569
- BROWN A AND PACHECO S (2006) Propuesta de actualización del mapa ecorregional de la Argentina. Pp. 25-31 in: Brown AU, Martínez Ortiz MAC, Pacheco J, editors. La Situación Ambiental Argentina 2005. Fundación Vida Silvestre Argentina, Buenos Aires
- CABRERA AL (1971) Fitogeografía de la república Argentina. *Boletín de la Sociedad Argentina de Botánica* 14:1-42
- FERNÁNDEZ FJ, BALLEJO F, MOREIRA GJ, TONNI EP AND DE SANTIS LJM (2011) Roedores cricétidos de la Provincia de Mendoza: Guía cráneo-dentaria orientada para su aplicación en estudios zooarqueológicos, Universitas, Córdoba
- FERNÁNDEZ F, DEL PAPA LM, MANGE E, TETA P, CRIVELLI MONTERO E, PARDIÑAS UFJ (2016a) Human subsistence and environmental stability during the last 2200 years in Epullán Chica cave (northwestern Patagonia, Argentina): A perspective from the zooarchaeological record. *Quaternary International* 391: 38-50
- FERNÁNDEZ FJ, MONTALVO CI, FERNÁNDEZ JALVO Y, ANDREWS P AND LÓPEZ JM (2017) A re-evaluation of the taphonomic methodology for the study of small mammal fossil assemblages of South America. *Quaternary Science Reviews* 155:37-49
- FERNÁNDEZ FJ AND PARDIÑAS UFJ (2018) Small mammals taphonomy and environmental evolution during Late Pleistocene-Holocene in Monte Desert: The evidence of Gruta del Indio (central west Argentina). *Journal of South American Earth Sciences* 84:266-275
- FERNÁNDEZ FJ, TETA P AND BARBERENA R (2012) Small mammal remains from Cueva Huemul, northern Patagonia, Argentina: *Quaternary International* 278:22-31
- FERNÁNDEZ FJ, TETA P, MANGE E, VENANZI LG AND PARDIÑAS UFJ (2016b) Micromamíferos del sitio arqueológico Cueva Galpón (Río Negro, Argentina): Aspectos tafonómicos y reconstrucción paleoambiental para el Holoceno Tardío en Norpatagonia oriental. *Arqueología* 5159:105-124
- JAYAT P, TETA P, OJEDA AA, STEPPAN SJ, OSLAND JM, ORTIZ PE, NOVILLO A, LANZONE C AND OJEDARA (2021) The *Phyllotis xanthopygus* complex (Rodentia, Cricetidae) in central Andes, systematics and description of a new species. *Zoologica Scripta* 50:689-706

- JAYAT P, TETA P, OJEDA AA, STEPPAN SJ, OSLAND JM, ORTIZ PE, NOVILLO A, LANZONE C AND OJEDA RA (2022) Establishing the availability of the recently erected binomen *Phyllotis pehuenche* (Rodentia, Cricetidae, Sigmodontinae). *Therya notes* 3:110-114
- LÓPEZ JM (2020) Actualistic taphonomy of barn owl pellet-derived small mammal bone accumulations in arid environments of South America. *Journal of Quaternary Science* 35:1057-1069
- LÓPEZ JM AND CHIAVAZZA H (2019) Small mammal bone remains recovered in open-air archaeological sites from the Monte Desert in Central Western Argentina: taphonomic and palaeoenvironmental implications. *Archaeological and Anthropological Sciences* 11:5149-5169
- LÓPEZ JM, AGUILAR JP AND FERNÁNDEZ FJ (2021a) Egagrópilas actuales de aves rapaces como herramientas para la generación de modelos paleoambientales en la provincia de Mendoza, Argentina. *Mastozoología Neotropical* 28:1-29
- LÓPEZ JM, QUINTANA MF, MIGNINO J AND PEREYRA-LOBOS R (2021b) Tafonomía de restos óseos y dentarios de micromamíferos recuperados en egagrópilas actuales generadas por la Lechucita de las Vizcacheras (*Athene cucularia*): una discusión sobre las tierras bajas áridas del Centro Occidente Argentino. *Intersecciones en Antropología* 22:55-67
- LYMAN R (2008) Quantitative paleozoology. Cambridge University Press, New York
- MARES MA (1992) Neotropical mammals and the myth of Amazonian biodiversity. *Science* 255:976-979
- MARES MA AND OJEDA RA (1982) Patterns of diversity and adaptation in South American hystricognath rodents in: Mares MA, Genoways HH, editors. *Mammalian Biology in South America*. Special Publication Series No. 6, Pymatuning Lab. Ecology: Linnesville (USA), p. 393-432
- MARTINEZ GN (2018) Nueva campaña GAEMN-GEMA en la Caverna del León, Neuquén. *EspeleoAr* 18:3-4
- MIGNINO J, IZETA A, GARCÍA JM, CATTÁNEO R AND PEDETTI OB (2021). Neo-tafonomía de restos de pequeños mamíferos acumulados por la Lechuza del Campanario (*Tyto alba*) en el Gran Chaco Sudamericano (Argentina): Un marco de referencia para sitios de percha y anidamiento. *El Hornero* 36:51-60
- MONTALVO CI AND FERNÁNDEZ FJ (2019) Review of the actualistic taphonomy of small mammals ingested by South American predators: its importance in the interpretation of the fossil record. *Publicación Electrónica de la Asociación Paleontológica Argentina* 19:18-46
- MONTALVO CI, FERNÁNDEZ FJ AND TALLADE PO (2016) The role of *Bubo virginianus magellanicus* as rodent bone accumulator in Archaeological Sites: A Case Study for the Atuel River (Mendoza, Argentina). *International Journal of Osteoarchaeology* 26:974-986
- MORELLO J (1958) La provincia fitogeográfica del Monte. *Opera Lilloana* II 2:5-155
- MURRIE M AND MURRIE S (2010) The First Book of Seconds: 220 of the Most Random, Remarkable, Respectable, Regrettable Runners-Up and Their Almost Claim to Fame. Simon and Schust. Avon (USA), 240 p
- OJEDA RA, BLENDINGER PG AND BRANDL R (2000) Mammals in South American drylands: Faunal similarity and trophic structure. *Global Ecology and Biogeography* 9:115-123
- OJEDA RA AND TABENI S (2009) The mammals of the Monte Desert revisited. *Journal of Arid Environments* 73:173-181
- PARDIÑAS UFJ (1999) Tafonomía de microvertebrados en yacimientos arqueológicos de Patagonia. *Arqueología* 9:265-340
- PARDIÑAS UFJ, UDRIZAR-SAUTHIER DE AND TETA P (2012) Micromammal diversity loss in central-eastern Patagonia over the last 400 years. *Journal of Arid Environments* 85:71-75
- PARUELO JM, BELTRÁN A, JOBBÁGY E, SALA OE AND GOLLUSCIO RA (1998) The climate of Patagonia: general patterns and controls on biotic processes. *Ecología Austral* 8:85-101
- PATTON JL, PARDIÑAS UFJ AND D'ELÍA G (2015) *Mammals of South America, Volume 2: Rodents*. London (UK): University of Chicago Press
- R DEVELOPMENT CORE TEAM (2020) R: A Language and Environment for Statistical Computing. R Foundation for Statistical Computing Vienna Austria
- SAAVEDRA B AND SIMONETTI JA (1998) Small Mammal Taphonomy: Intraspecific Bone Assemblage Comparison between South and North American Barn Owl, *Tyto alba*, Populations. *Journal of Archaeological Science* 25:165-170
- TARQUINO-CARBONELL AP AND TURIENZO P (2019) Presencia de *Histiopus montanus* (Chiroptera: Vespertilionidae) en la Caverna del León (Neuquén, Argentina). *Gota a Gota* 17:44-49